



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

COMMUNICATIONS FROM THE HARVARD MINERALOGICAL  
MUSEUM.

OCCURRENCE OF NATIVE COPPER AT FRANKLIN  
FURNACE, NEW JERSEY.

By J. E. WOLFF.

Presented April 13, 1898.

In 1897 a large specimen of ore containing native copper from the new (Parker) shaft at North Mine Hill, Franklin Furnace, was acquired for the Harvard Mineralogical Museum.

The specimen, some seven inches square, is broken flat parallel to the banding of the ore body, and is composed of a central band of apple-green willemite interbanded with coarse white calcite and smaller masses of franklinite and zincite in the usual association. The cleavage surfaces of the calcite masses are curved by pressure, and show the development of the gliding planes parallel to  $-\frac{1}{2} R$ . In the central zone of the specimen these evidences of pressure become intensified, and small slickensided surfaces are developed in the willemite and calcite, while the former mineral is irregularly cracked in addition to the development of the imperfect cleavage. In this zone native copper occurs, forming a band half an inch wide running parallel to the banding of the ore. The copper has no distinct crystalline form, but occurs as thin films or thicker sheets filling the cracks in the willemite and occasionally in the calcite. Except for traces of oxidation on its surface, the copper is pure and associated with no other mineral.

Of two other specimens loaned by Mr. J. A. Van Mater, superintendent at North Mine Hill, one has the same association described above, and shows a veinlet of copper penetrating calcite along a gliding plane, in addition to the penetration along fissures in the willemite. In the other the copper occurs in willemite which is associated with yellow garnet, black biotite, and calcite. The masses of copper are thicker here, and a projecting point laid bare by the removal of the matrix has some resemblance to a spear-head twin of copper, but is probably a matrix mould.

In May, 1898, small sheets of native copper were seen by the writer in vein matter coming up from the deep levels, but have not yet been observed in the standing vein. The matrix is a light yellow densely granular mineral with resinous lustre, which appears from preliminary tests to be a phase of the yellow manganese garnet (polyadelphite) so common in the vein matter of the Parker shaft. It is intimately mixed in microscopic veinlets and larger masses with an amorphous (often earthy) light pink manganese mineral not further identified. A thin section of this gangue shows cloudy yellow spots of the garnet surrounded by paler transparent material which has the feeble double refraction in sectors of some garnet, a property which has not been observed by the writer in the ordinary deep yellow garnet of the ore body. This abnormal character and the veining by the pink material suggest an alteration of the original garnet gangue with which the deposition of the copper might have been contemporaneous.

Mr. Van Mater states that native copper occurred in 1897 about eight hundred feet below the surface in a seam in the middle of the ore body, i. e. about half way between the hanging and foot walls. This seam was filled with a gangue of rhodonite and other minerals in an earthy porous condition, and probably represents the form just described.

The other occurrence in massive willemite was not observed in the standing vein, but was picked out from the ore coming from about the same depth.

It is evident, from the relations of the veinlets of the copper to the cleavage and cracking of the willemite and secondary pressure planes of the calcite, that its deposition took place *after* the formation of the zinc ores and associated minerals, and that possibly the circulation of the solutions from which it was deposited took place along the same motion planes in the ore body which produced the beautiful slickensides found occasionally in this part of the deposit. There are no facts on which to base a conclusion as to the form in which the copper was carried in solution, nor as to the method of its final reduction and deposition, although it is probable the changes noticed above in the garnet gangue were connected with it.

This occurrence increases the long list of minerals found at Franklin Furnace and vicinity (now nearly seventy), and is an interesting addition to the occurrences of native copper below the zone of surface oxidation.